


# Protection of Wild Adult Steelhead in Idaho by Adipose Fin Removal

Final Report  
1985 - 1988



This Document should be cited as follows:

*Duke, Rodney, "Protection of Wild Adult Steelhead in Idaho by Adipose Fin Removal", Project No. I 984-00200,, 4 7 electronic pages., (BP4 4 Report DOE/BP-I4903-)*

Bonneville Power Administration  
P.O. Box 3621  
Portland, Oregon 97208

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

PROTECTION OF WILD ADULT STEELHEAD IN IDAHO  
BY ADIPOSE FIN REMOVAL

Final Report

By

Rodney Duke, Project Leader

Idaho Department of Fish and Game  
1540 Warner Avenue  
Lewiston, Idaho 83501

Funded by:

Ronald Morinaka, Project Manager  
U.S. Department of Energy  
Bonneville Power Administration  
Division of Fish and Wildlife  
P.O. Box 3621  
Portland, Oregon 97208

Projects 84-2  
Contract No. DE-A-179-84BP14903

April 1989

## TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT .....	1
INTRODUCTION .....	2
OBJECTIVES .....	2
METHODS .....	2
RESULTS .....	4
1987-1988 Mark Operations .....	4
1983-1988 Mark Operations .....	6
1985 Clip Healing Studies .....	6
Adipose Fin Histological Examination .....	6
Returning Adult Clip Quality. ....	6
Hatchery Versus Wild Adult Returns .....	8
DISCUSSION .....	8
Hark Quality .....	8
Effect of the Program on Wild Runs .....	9
Clip Healing .....	10
Adipose Fin Composition and Purpose .....	10
Fin Regeneration .....	11
Marked Fish Survival .....	12
CONCLUSIONS .....	13
BIBLIOGRAPHIES .....	15
ACKNOWLEDGEMENTS.....	16
LITERATURE CITED .....	17
APPENDICES .....	19

## LIST OF TABLES

Table 1.	Summary of Idaho hatchery steelhead adipose fin marking operations, 1987 .....	5
Table 2.	Five-year adipose marking summary for Idaho hatcheries, 1983-1985 .....	7

## LIST OF APPENDICES

	<u>Page</u>
Appendix A. Bibliography - marked fish survival .....	20
Appendix B. Bibliography - fin regeneration .....	26
Appendix C. Bibliography - hooking mortality .....	32



## ABSTRACT

Steelhead trout (Salmo gairdneri) reared in Idaho hatcheries for release during the 1984-1988 outmigrations were adipose fin clipped to differentiate between wild/natural and hatchery-reared fish. From 1984- 1988, 34 million hatchery-reared steelhead trout were clipped and 30.1 million were released; the difference being made up by hatchery mortality and the percent of acceptable clips. Since 1987, the adipose clip has given protection to all wild/natural steelhead and identified them from hatchery stocks.

It is premature to conclude whether adipose marking can be totally effective in preserving wild runs into Idaho. Although wild runs appear to be increasing, the full effect of this program cannot be evaluated until the mid-1990s, when sufficient numbers of year classes will have returned to provide significant data. In the interim, the program is successfully protecting wild stocks from being harvested in Idaho's sport fishery. This program, if continued, will undoubtedly reduce harvest mortality of wild stocks and enhance their restoration.

Author:

Rodney C. Duke  
Senior Fishery Research Biologist

## INTRODUCTION

In the early 1980s, the improved survival of hatchery-reared steelhead created stock management problems. While adequate numbers of hatchery-reared fish led to harvestable surpluses, wild stocks remained at low levels. Angler harvest directed at hatchery stocks led to overharvest of wild stocks. In order to harvest surplus hatchery stocks and protect wild stocks in the fishery, anglers needed a means of identifying between the two at the time the fish were caught. Initially, various dorsal fin length criteria were used to regulate harvest. These were not totally successful because of variability and enforcement difficulties. To solve the problems associated with differential harvest, fishery scientists decided to desquester the adipose fin clip previously used to identify the presence of a coded wire tag and use it as a harvest identifier. All hatchery-reared steelhead in the Columbia River Basin were to be adipose clipped.

On November 1, 1983, the Idaho Department of Fish and Game secured a five-year contract from Bonneville Power Administration (BPA) to adipose clip all of the hatchery-reared steelhead in Idaho to aid in the protection of Idaho's wild stocks. This report summarizes that work.

## OBJECTIVES

The objectives of the project were:

1. To remove 751 of the adipose fins from 99% of the hatchery-reared steelhead in Idaho.
2. To review current information on the use of adipose fin clipping.
3. To determine, if possible, the physiological role of the adipose fin.

## METHODS

From 1983 to 1987, fish marking operations were conducted at three Idaho steelhead production facilities: Niagara Springs (Idaho Power), Hagerman National, and Dworshak National fish hatcheries. In 1987, the Magic Valley Fish Hatchery was put into production, and all fish were marked for release in the spring of 1988.

Except for the first year, all marking was conducted in two specially built mobile marking units. These units were constructed in 1984 specifically for the large marking operations previously handled by various departmental and federal marking trailers. The new marking units each contain 16 work stations and efficiently handle large numbers of fish with minimal stress (Duke 1985).



Inside the trailers, fish were placed in troughs containing a solution of MS-222, Propolyaqua, and anti-foam. The solution was recirculated, cooled, and aerated. MS-222 is an anesthetic, Propolyaqua is a stress reducer, and anti-foam prolongs the effect of the anesthetic solution. Once the fish are anesthetized, personnel remove the adipose fin with scissors.

Quality inspections were taken throughout the marking operation to ensure that the clip was being done properly and that all fish were clipped. Prior to release, a minimum 300 fish were visually examined to determine the quality of the clip. Examined fish were divided into five categories based on their fin clip: excellent (100% removal); too deep (100% removal, but secondary infections or complications possible); poor (less than 10% remaining at the leading edge); partial (10-25% remaining); and not clipped (more than 25% remaining). All clip categories, except "not clipped," are acceptable to be used in identifying returning adult fishes.

Fish health was monitored throughout the rearing cycle and mortalities noted. Mortality attributable to marking was calculated by subtracting the daily mortality before marking began from the mortality level after marking until the mortality rate stabilized at pre-marking levels. In some cases, rearing histories from the previous year were used to estimate the normal daily mortality rate for that particular stage of rearing.

In 1985, in vivo experiments on healing rate were conducted at Dworshak National Fish Hatchery. Both unclipped (control) and clipped (test) fish were held in circular vats. Idaho Department of Fish and Game personnel monitored the tests and documented the healing process photographically. Samples of juvenile and adult adipose fins were sent to the U.S. Fish and Wildlife Service's Cultural Development Center where Charlie E. Smith examined histological sections for tissue composition.

Extensive reviews of literature on hooking mortality rates, the use of the adipose fin marking system in the protection of wild fish, adipose fin function and composition, and effects of fin clipping on survival were gathered from various sources and were included in a previous annual report (Duke 1986). They have also been included at the end of this report.

In 1986, fish from the first ad-clip marking operation returned to Idaho. Returning one-salt adult steelhead were monitored to determine the quality and degree of regeneration of the adipose clip. In the fall of 1987 and spring of 1988, a more extensive check of adult clip quality was conducted in both the sport fishery and at the spawning racks. Hatchery and wild compositions were determined as the fish passed through various adult monitoring facilities located at selected hydroelectric projects. This was the first year that all fish in the three returning age classes were adipose clipped under this marking program.

## RESULTS

### 1987-1988 Mark Operations

Clipping for the final year of this project began on September 8, 1987 at Niagara Springs Steelhead Hatchery and continued until October 1, 1987. A total of 2,300,818 hatchery steelhead were clipped. The marking operation was then moved to Hagerman National Fish Hatchery from October 5 until October 23, 1987, where 1,986,060 steelhead were clipped. Steelhead were clipped at the Magic Valley Steelhead Hatchery during the period October 27 through November 20, 1987. A total of 2,116,071 steelhead were marked at Magic Valley.

The same marking personnel were used at all three facilities. Crews worked two 6-hour shifts in a single mobile marking unit. A total of 554 hours (about 47 days) were required to mark the 6.4 million fish at the three hatcheries, representing an average of 11,645 fish clipped per hour at the three facilities.

Due to the number of fish marked at Hagerman, Niagara, and Magic Valley, marking at Dworshak was delayed until December 1987. Consequently, Dworshak's fish nearly exceeded the optimum marking size. This, coupled with the large number of fish at Dworshak, presented major challenges in order to complete the operation before the January 1 cutoff date for marking at the facility. In order to complete marking prior to January 1, two adipose marking trailers were used in conjunction with the two coded wire tagging (CWT) trailers. Those fish requiring coded wire tags were adipose clipped during the CWT operation to prevent double handling of fish. After each CWT group was completed, the remaining fish in a pond were adipose clipped. Ten days were required to adipose clip 2,973,776 fish at Dworshak. Over 130 different temporaries were hired. Two 8-hour shifts were run in each trailer, requiring two supervisors for each shift. Because of mechanical problems, not all trailers were continually in operation. Extensive moving of the mobile units was also necessary. The operation, however, was highly successful, and an average of 297,377 fish were clipped (sometimes tagged and left-ventral clipped as well) per day. This translates to 24,781 fish per hour being marked.

In general, fish at all four facilities appeared healthy and in good condition. However, viral disease was confirmed at Niagara and Dworshak during marking. Fish in infected raceways were marked last at both facilities to prevent the spread of disease. Initial marking mortality at all facilities was 0.0632 of the total fish marked. Initial clip quality checks at the southern hatcheries indicated less than 12 were not clipped. Visual quality checks were taken at Dworshak because of time limitations. Sampling for quality checks was done prior to release. Table 1 summarizes the results of the quality checks from the four facilities, as well as the 1987 clipping operations.

Table 1. Summary of Idaho hatchery steelhead adipose fin marking operations, 1987.

	Niagara Springs Hatchery	Hagerman National Hatchery	Magic Valley Hatchery	Dworshak National Hatchery	Total
No. of fish marked	2,300,818	1,986,060	2,116,071	2,973,725	9,376,725
No. of days required	18	13.5	15.33	10	56.83
No. of man hours required <sup>a</sup>	3,456	2,592	2,943	3,096	12,087
No. of fish clipped/man hour <sup>b</sup>	665.7	766.2	719.0	950.5	
Initial clipping mortality	1,711	941	883	2,685	6,220
Percent mortality	.074	.047	.042	.090	.063
Percent acceptable clzps <sup>c</sup>	99.76	99.90	99.88	97.79	99.85

<sup>a</sup>Markers only.

<sup>b</sup>At Niagara Springs, Hagerman Natl., and Magic Valley, two 6-hour shifts were operated out of one marking unit. At Dworshak Natl., two 8-hour shifts were operated out of two ad-marking units. Additional fish were marked in conjunction with tagging operations in two CWT trailers.

<sup>c</sup>Quality checks are initially taken during the mark operation, except for Dworshak, which is a final quality check at release.

### 1983-1988 Mark Operations

During the five years of the project, nearly 34 million steelhead trout were adipose clipped. Final hatchery releases of clipped fish totaled 30,121,306. During this period, BPA contract expenses totaled \$441,000. The total cost of adipose clipping (based on total hatchery release) over the five-year contract was \$14.65/thousand. Table 2 summarizes the five years of marking operations.

### 1985 Clip Healing Studies

Observations of fish in the raceways at all hatcheries indicated complete healing of the tissue in the area of excision within 3 to 4 weeks. Fish that had been clipped too deeply usually showed signs of fungus within one week. There were also observed cases of nipping at the excised white area. In severe cases, fish with fungus died and accounted for the largest percentage of the observed delayed mortality. Mortality from this cause ended within four weeks. Observation from the in vivo test at Dworshak National Fish Hatchery indicated healing was rapid and complete within 22 days (Pat Chapman, IDFG, personal communication). From observation and photographs, there was no apparent difference in the various full and partial clips in terms of regeneration or healing.

### Adipose Fin Histological Examination

Histological analysis for both juvenile and adult steelhead trout showed the adipose fins to be composed primarily of fibrous connective tissue interspersed with few blood vessels and occasional adipose (fat) cells. Special connective tissue stain demonstrated a high percentage of collagenous fibers typical of fibrous connective tissue. There was no evidence of glandular or secretory epithelial cells which could be associated with hormonal production. There were no differences in the composition of fins from juvenile and adult steelhead trout (Charlie Smith, personal communication).

### Returning Adult Clip Quality

Adult steelhead returning to various hatchery spawning racks during 1986 and 1987 were examined for adipose fin clips. Only those fish of obvious hatchery origin (i.e., irregular fin rays) were included in the samples. At Dworshak in 1986, 33 out of 701 one-salt fish examined had adipose fins, or 4.72. In 1987, 13 out of the 293 examined (4.4%) of the one-salt fish had adipose fins. An examination of 4,869 two-salt fish in 1987 showed 196 with adipose fins for a 4.02 adipose fin retention. In 1988, 3,784 adult steelhead were examined, and 152, or 4.0%, were nonadipose clipped. An age class breakdown was not available for the 1988 returning fish.

Table 2. Five-year adipose marking summary for Idaho hatcheries, 1983-1985.

	1984	1985	1 9 8 6	1987	1988
Hagerman National					
No. fish clipped	NA <sup>b</sup>	1,421,6	1,836,058	1,777,997	1,986,060
x acceptable clips		99.30	99.70	99.35	99.90
Total hatchery mortality		— <sup>c</sup>	183,137	256,372	274,123
Total hatchery release	1,162,475	1,526,932	1,652,921	1,521,625	1,711,937
Niagara Springs					
No. fish clipped	NA	2,502,125	2,291,784	1,928,982	2,300,818
X acceptable clips		99.70	99.70	99.30	99.76
Total hatchery mortality		483,592	611,931	77,082	511,792
Total hatchery release	1,822,805	2,018,533	1,679,853	1,851,900	1,789,026
Dworshak National					
No. fish clipped	NA	2,436,7	3,261,429	1,955,144	2,973,725
X acceptable clips		99.30	99.33	99.36	97.79
Total hatchery mortality		835,028	312,952	--~	271,498
Total hatchery release	1,961,392	1,601,6 05	2,948,477	2,105,748	2,702,227
Magic Valley					
No. fish clipped	--	-- --		--	2,116,071
X acceptable clips	- - -	- - -	- - - (NEW FACILITY)	-----	99.88
Total hatchery mortality					52,311
Total hatchery release					2,063,760
Year total (hatchery release)	4,946,672	5,147,160	6,281,251	5,479,273	8,266,950
Grand total (hatchery release)	30,121,306				

<sup>a</sup>Total hatchery mortality includes all mortality monitored from time of marking to release. All post-marking mortality may not be directly related to the mark operation.

<sup>b</sup>Mark data and mortality not available because of multiagency involvement and program was not in place at time of marking. All steelhead in Idaho were clipped prior to release.

<sup>c</sup>Difference due to disagreement in either count and/or hatchery inventory. All fish on-station were clipped.

uality check samples were taken at the Pahsimeroi, East Fork, Sawtooth, and Oxbow return racks in 1988. These fish were released from Hagerman National and Niagara Springs hatcheries. At the Pahsimeroi facility, 1,094 fish were checked, and 8 had unacceptable clips, for a 0.73% adipose fin retention. At Sawtooth, 837 adult steelhead were examined, and 5 bad clips (62) were found. At the East Fork trap, 190 fish were examined, and none were found with adipose fins. Adult steelhead were examined at Oxbow Hatchery, and 43 out of 2,524 (1.7%) were found to be unclipped hatchery-reared steelhead. Fifty-six fish (4.32) were identified as wild/natural at Oxbow. The relatively higher numbers of unclipped fish may have resulted because of straying from Oregon hatcheries, where not all returning hatchery fish had been clipped (Rich Carmichael, personal communication). Analysis of mark data to determine the actual percentage of adipose fin retention on Idaho stocks will not be available until late 1988. In 1986, 0.32 of the returning adult fish at the Pahsimeroi still had intact adipose fins. Data is not available for the other sites in 1986 or any site in 1987.

### Hatchery Versus Wild Adult Returns

A complete analysis of the present and future status of wild and hatchery steelhead trout in Idaho is beyond the scope of this project. Personnel from various state and federal agencies are involved with monitoring both run segments as they enter the Columbia River and migrate into Idaho and through the numerous fisheries. The percent composition of hatchery and wild fish at Lower Granite Dam, based upon dorsal fin erosion, was determined in 1984. That fish run had no clipped fish from this project but was 242 wild. The next year, 1985, scale analysis showed 23% of the fish were wild. In 1986, scale analysis showed the wild run segment at 20.4% of the total run.

## DISCUSSION

### Mark Quality

The experiences during the past five years demonstrate that an entire state's steelhead production can be marked in an efficient manner. Quality checks on adults returning from fish released from southern Idaho facilities consistently indicate about 1% of the returning adults have adipose fins that were missed in the fin clip operation. However, at Dworshak, the percent of missed adipose clips is higher (near 4%).

The total number of fish unclipped at Dworshak is partly due to crowding problems during mark operations which are associated with design problems with the Burroughs ponds. It is difficult, if not impossible, to remove every fish from the ponds because of the rounded corners and fill pipes located at the outer corners. Extra care is taken, but smaller fish, in particular, can elude the crowd rack by going among the pipes. Additional complicating factors include: 1) Fish are often larger than

those at other facilities at the time of marking. The larger size makes it difficult for the markers to handle the fish, and a few can slip from the marker's hands and escape into the return pipe inside of the trailer without being clipped. 2) There is usually a wide size variation in the fish. In some years, the percentage of "pin heads" is high in comparison to the southern facilities because "drop out" usually has not occurred. The adipose fin is easily missed on these fish (less than 60 mm in total length) when marked with fish exceeding 150 mm. 3) Crowd screens must be of a large enough mesh to prevent clogging during marking operations. However, this size also allows some pin heads to escape and go unclipped. 4) Fish escape to the river in normal hatchery operations prior to marking. This problem occurs at all facilities to a minor degree during the course of normal hatchery operations. Fish escaping at the southern facilities are prevented from migrating to the ocean because of several migrational barriers. Only those fish escaping at Dworshak have the potential of migrating and entering the fishery.

The number of unmarked fish due to any of the above factors is variable. Those factors associated with "pin heading" are quite minor since there are indications these fish do not survive to migration. However, the combination of operational errors inside the marking units and at the hatchery due to either handling or design can result in missed fish. It may be difficult to ever obtain better than a 96% adipose-clipped population at Dworshak. Although the percentage of missed clips at Dworshak is almost four times higher than other facilities, it does not present serious problems to the management and protection of the wild runs into the Clearwater River. The total number of harvestable hatchery fish is relatively low.

#### Effect of the Program on Wild Runs

It is premature to conclude the degree of effectiveness of the adipose marking program in preserving wild runs into Idaho. The wild runs seem to be increasing as determined by the percent composition of the runs, especially for the 1987-1988 fish run. However, the increase in the wild/natural population is probably a function of several programs giving results. The limited protection given wild adults by the dorsal fin regulation, adult outplantings, and fry plantings are all contributing to a wild/natural comeback. Certainly the adipose marking program is protecting wild stocks in Idaho by eliminating excessive harvest, especially in the early part of the season when wild fish are most vulnerable in areas of high fishing pressure. McArthur (1988), in a telephone survey he conducted statewide, estimated that during the 1987 steelhead season, there were 13,152 wild fish released as a direct result of the adipose fin regulation. Though some fish could have been caught and released more than once, the impact of the adipose fin regulation initiated by this project will undoubtedly have a major positive impact on Idaho's wild steelhead runs in the future. But because of the life history of the fish, it will be 1995 before the full impact of this program can be evaluated.

### Clip Healing

Healing of the area of excision is dependent on several factors, including fish health, water quality and temperature, and size at clipping. Larger fish tend to expose more tissue (white area) and usually suffer higher losses due to nipping, which turns into "sore back." Less "sore back" is observed when fish are clipped at smaller sizes. Clips which are too deep expose even more tissue area and result in severe cases of nipping and "sore back." Secondary infections, such as fungus, also result from too deep a clip. In these cases, the fish usually never heals, and death occurs within one month.

Water temperature is critical and can be beneficial or detrimental. As a rule, fish tend to heal faster in warmer water (10°F), but fungus and secondary infections usually increase. Conversely, fish in cold water (1 to 5°C) heal more slowly but show less signs of fungus and secondary infections. Fish marked in cold water temperatures appear to take the stress of the handling operation better, and fewer mortalities result.

In general, fish heal remarkably fast, and from observed and in vivo observations, healing time is not a critical factor in the marking operation. Post-marking prophylactic treatments with quaternary ammonia compounds usually control any complication resulting from the excision of the adipose fin.

### Adipose Fin Composition and Purpose

The salmonid adipose fin is relatively small, fleshy, and immobile. It does not contain any skeletal elements, is scaleless, and covered only by the dermis and epidermis. It is filled with an amorphous matrix of loose connective tissue (Harder 1975).

Historically, the adipose fin was thought to be a vestigial fin without purpose and comprised mostly of fatty tissue and thus the name adipose. Weisel (1968) found no fat in sockeye salmon (Oncorhynchus nerka) and only 3 to 4 drops section in the cutthroat trout (Salmo clarki) and brown trout (Salmo trutta). In recent years, there have been several hypotheses that the fin may serve other functions, ranging from hormonal regulation to pH balance. No literature to substantiate any of these hypotheses could be found. The histological examination found no evidence of glandular or secretory epithelial cells which normally are associated with endocrinal or hormonal functions. Stuart (1958) does make mention that the size and shape of the adipose fin is a secondary sex characteristic in spawning salmonids, with the female fin long, thin, and narrow at the base in comparison to a short, thick, wide-based male fin.

Aleyev (1977) suggested the adipose fin in salmonids functioned basically the same as finlets in the Scombridae (mackerel, tuna, etc.), i.e., it functions to transversely streamline the caudal peduncle for faster swimming.



## Fin Regeneration

The practice of fin clipping fish for the purpose of identification has been used for many years. There are many advantages and disadvantages of this type of identification procedure. A question in all studies utilizing fin clipping is whether the fin will partially or completely regenerate, thus rendering the mark difficult to recognize at a later date. Several studies have been conducted to evaluate the regeneration of the various types of fins.

Mears (1976), studying brook trout (Salvelinus fontinalis), found regeneration had occurred in 92% of all fins observed. The frequency of regeneration was highest (41%) for the anal fin and lowest (9%) for the adipose. Regeneration of the pectoral fins was 1.5 to 2.0 times more frequent than that of pelvic fins. Few fins regenerated to more than 50% of their original size.

Johnson and Shelton (1958) found little regeneration of either adipose or pectoral fins on fall chinook at the Spring Creek Hatchery, with 99.2% of all fish with a double fin mark easily identified as double-marked fish four months after marking.

Stauffer and Hansen (1969) utilized left maxillary, right maxillary, adipose, right pectoral, and right pelvic clips to mark rainbow trout. After two years in the hatchery, 95% or more of the pelvic, pectoral, and maxillary clips were recognizable (one-half or less of the fin regenerated). There was no regeneration of the adipose fin.

Shetter (1951) marked four groups of lake trout (Salvelinus namaycush) with various fin clips. The adipose clip had 5.2% regeneration, the dorsal 6.4%, right pectoral 3.5%, left pectoral 10.2%, and right ventral 35.9% (one-half or more of the fin remaining). However, there appears to be some ambiguity in his reference to the quality of the clipping operation, especially for the right ventral clip.

Hale (1954) found pelvic fin regeneration to be "complete" (both fins similar and normal in size and spread) in 31.5% of the brook trout he held for 14 months 10 days. Most of the remaining fish showed "partial" (one fin club-like in appearance and the other club-like or completely regenerated) or "considerable" (one or both fins somewhat smaller and spreading like a normal fin) fin regeneration (13.7% and 47.5%, respectively). He attributed this high regeneration to inexperience and lack of skill of those clipping small fish (3.0 inches total length).

Other authors have also reported varying results. However, the various studies indicate that the adipose has the least regeneration, with the pelvics next, followed by the pectorals and anal, in respective order. In each study reviewed, a hypothesis was made that a fin with 50% remaining was regenerated. Few studies acknowledged that the regeneration could be aggravated by poor clipping technique. For many studies, the percentage of regeneration may be nothing more than an indication of initial clip quality. Experience and results of marking juvenile salmon

and steelhead for the IDFG coded-wire tagging program has shown that adipose fins properly and totally excised will not regenerate.

Examination of adult steelhead and salmon marked with an adipose clip and coded-wire tag as a juvenile indicates there is no regeneration in a complete clip and only slight regeneration of a partially clipped fin as the incision heals. However, these are usually recognizable as a clip and only when approximately 25% or less of the fin is removed does the mark become questionable. Fish marked with a coded-wire tag and left ventral clip indicate no regeneration of the clip if the fin is clipped below the basal bone. However, leaving only a few fin rays can result in an entire, but usually deformed, fin. Again, the amount of regeneration is directly related to the quality of the clip. In the case of the ventral clip, a poorly applied clip usually results in the loss of information since it is difficult to differentiate a regenerated fin on a hatchery fish from the deformation the fin may have received during rearing.

Stuart (1957) details pictorially the regeneration of partially clipped fins. The observations made in the coded-wire tagging program are basically in agreement with his study.

#### Marked Fish Survival

Experimentation utilizing marked fish assumes no differential in survival of marked fish and their unmarked counterparts they represent. However, there are studies reporting differential survival of the various clips. A differential in survival may be a result of interference with swimming ability, endurance, behavior, or growth. Other factors, such as handling during marking, physiological condition of the fish at marking, size at marking, and others, may also play an important role.

Bonham (1968) concluded that a maxillary mark on chinook salmon (Oncorhynchus tshawytscha) retarded growth and probably induced mortality and was less desirable than a ventral clip. Wales (1947) found the survival of pelvic-clipped fingerling brook, brawn, and rainbow trout to be better than similar lots of pectoral fin-clipped fish. Shetter (1951) tested the survival of lake trout receiving a single pectoral, single pelvic, or dorsal plus adipose fin clip. After correction for fin regeneration, he concluded there was no significant difference between the survival rates of the marked and unmarked fish. Armstrong (1949) found no difference in the survival rates of unmarked and adipose-clipped lake trout fingerlings after ten months. Experiments conducted by Nelson (1960) indicated that the removal of the pelvic, adipose, or dorsal fins did not significantly affect the survival of fingerling brook or rainbow trout under hatchery conditions. Other authors, such as Ricker (1949) and Foerster (1936), found that marked fish survived significantly less than unmarked fish during their studies. In another experiment, Shetter (1952) found that fin-clipped fingerling lake trout (utilizing the same clips as in his previous work) did not suffer any greater losses from predation than did unmarked fish. Horack (1969) using a stamina tunnel tested the swimming ability of 3.3- to 4.0-inch rainbow trout. He found that swimming ability was not significantly affected by removing either the

dorsal, both pelvic, both pectoral, anal, or adipose fins. He concluded that the removal of both ventral or pectoral fins may reduce stamina and should be used with caution. A caudal clip severely reduced stamina and could affect the outcome of studies. Nicola and Cordone (1973) studied the long-term survival of fin-clipped and unmarked rainbow trout in Castle Lake, California. They found that any fin clip was detrimental. The adipose fin clip reduced survival by as much as 50%. The removal of a ventral fin reduced survival by as much as 60% to 70%. The removal of a pectoral or dorsal fin reduced survival 70% to 80%. The removal of an anal fin was not any worse than the removal of the pectoral or dorsal, but removal had an inconsistent effect. They concluded that the anal fin clip should be avoided unless full excision could be guaranteed.

In the preliminary analysis of the Oregon Coastal Zone Management Association (OCZMA) proposal to mark hatchery-reared coho in the Oregon Production Index (OPI) area in 1984, a special task team evaluated the effects of marking. They concluded that in all experimental cases reviewed, extra handling and stress of fin clipping caused reduced survival. It also appeared that the survival effect went beyond immediate mortality due to stress in the hatchery. Although they could not identify the specific cause, evidence suggested that the removal of the adipose fin causes reduced ocean survival compared to unclipped fish. Loss of other fins or maxillary bones caused even greater losses. From the best available data, they concluded that adipose fin clipping would cause the loss of 52 to 202 of the hatchery coho production compared to unclipped releases. However, the data they present (Table 6, page 23 of the proposal), with the exception of one group, shows the adipose clip in conjunction with another mark. In almost all cases presented, the groups receiving a second mark in addition to the adipose clip survived less than the one group receiving only an adipose clip. The sources they used to estimate the 5% to 20% mortality for an adipose clip are not cited. Evidence from Idaho's fish marking, coded-wire tagging, and freeze branding operations indicates that mortality increases with each additional mark applied to the same fish.

Though there are many studies documenting marked fish survival and the differential survival between marked and unmarked fish, it appears from the literature available that the removal of the adipose fin affects survival of the fish the least, followed by the pelvics, pectorals, and dorsal. The caudal, anal, and maxillary are the least desirable to use, and studies indicate they may significantly affect survival.

## CONCLUSIONS

During the first five years of the marking program, personnel have been successful in achieving the marking of essentially all hatchery-reared steelhead in Idaho. During this time, 30,121,306 adipose-clipped steelhead trout have been released. Marking mortality during this time has been minimal, although total hatchery mortality prior to release in some situations is high, mainly because of viral diseases. Although wild runs appear to be increasing, the full effect of this program cannot be evaluated until the mid-1990s. However, in the interim,

the program is protecting wild stocks from being harvested in Idaho and allowing for successful harvest of hatchery stocks in a mixed-stock fishery.

The adipose fin mark is economically applied to large numbers of fish with minimal stress and mortality, affords a permanent and readily identifiable mark with good angler recognition, and provides for an easily enforced regulation. If all agencies and tribes in the Columbia Basin would participate fully by marking all hatchery-reared steelhead, wild/natural stocks could be protected in sport fisheries and more accurately accounted for in gill net fisheries. Adipose marking of hatchery steelhead will be continued in Idaho's fish hatcheries.

## BIBLIOGRAPHIES

The bibliographies presented are by no means complete. They are compilations from several research projects, books, and journals. Paul Mongillo (1984) and Victor A. Lewynsky (1980) both have done extensive reviews of literature on hooking mortality. Their work, along with others, is included.

## ACKNOWLEDGEMENTS

During the past five years, several department personnel have assisted in various phases of this project. Those permanently assigned to the project, and who have supervised marking operations and the writing of various reports, include: Arnie Miller, Todd Garlie, William Young, Timothy Wik, Bradford Dredge, and Doug Young. Bradford Dredge and Doug Young were in charge of marking operations during 1987. Joseph Norton and Rodney Duke assisted in the 1987 marking operation at Dworshak. Both Bradford Dredge and Joseph Norton assisted in preparation of this text.

## LITERATURE CITED

- Aleyev. 1977. *Nekon*. The Hague: Junk. (Reference translated and cited in R.W. Blake, 1983. *Fish Locomotion*).
- Armstrong, G.C. 1949. Mortality, rate of growth, and fin regeneration of marked and unmarked lake trout fingerlings at the Provincial Fish Hatchery, Port Arthur, Ontario. *Transactions of the American Fisheries Society* 77:129-131.
- Blake, R.W. 1983. *Fish Locomotion*. Cambridge University Press Cambridge, Great Britain.
- Bonham, K. 1968. Growth inhibition by maxillary excision, size at start of jaw elongation, and regression of weight on length of precocious male chinook salmon. *Transactions of the American Fisheries Society* 97(2):198-201.
- Duke, R.C. 1985. Final report, project AFS-1:Improvement of anadromous fish stocks and fisheries. Subproject 1, Study 1, Job 3: Steelhead marking trailers. Idaho Department of Fish and Game, Boise, Idaho.
- Duke, R.C. 1986. Protection of wild adult steelhead in Idaho by adipose fin removal. Annual report. U.S. Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife. Projects 84-2. Contract No. DE-A-179-84BP14903.
- Foerster, R.E. 1936. The return from the sea of sockeye salmon (*Oncorhynchus nerka*) with special reference to percentage survival, sex proportions and progress of migration. *Journal of the Biologist Board of Canada* 3(1):26-42.
- Hale, J.C. 1954. Regeneration of pelvic fins on fingerling brook trout following fin clipping. Investigation Report 1149, Minnesota Department of Conservation.
- Harder, W. 1975. *Anatomy of Fishes Part I*. Translated by Stephen Sokoloff, Tuebingen. E. Schweizerbart'sche Verlagsbuchhandlung (Magele V. Obermiller) Stuttgart.
- Horack, D.C. 1969. The effect of fin removal on stamina of hatchery-reared rainbow trout. *Progressive Fish-Culturist* 31(4):217-220.
- Hulbert, P.J., and R. Engstrom-Heg. 1980. Hooking mortality of worm-caught hatchery brown trout. *New York Fish and Game Journal* 27(1):1-10.
- Hunsaker, D., II, L.F. Marvell, and F.P. Sharpe. 1970. Hooking mortality of Yellowstone cutthroat trout. *Progressive Fish-Culturist* 32(4):231-235.

- Johnson, H.E. and J.M. Shelton. 1958. Marking chinook salmon fry. Progressive Fish-Culturist. 204(4):183-185.
- McArthur, T.J. 1988. Estimated 1987 spring steelhead season harvest and effort (telephone survey). Job Performance Report F-73-R-10, Subproject 2, Salmon and Steelhead Investigations, Study 1, Job 1. Idaho Department of Fish and Game.
- Mears, H.C. 1976. Overwinter regeneration of clipped fins in fingerling brook trout. The Progressive Fish-Culturist 38(2):73.
- Nelson, W.C. 1960. A comparison of the effects of the removal of the adipose and pelvic fins on fingerling brook and rainbow trout. Colorado Department of Game and Fish, Denver, Colorado.
- Nicola, S.J. and A.J. Cordone. 1973. Effects of fin removal on survival and growth of rainbow trout (Salmo gairdneri) in a natural environment. Transactions of the American Fisheries Society 102(4):753-759.
- Pettit, S.W. 1977. Steelhead catch-and-release research and regulations on the Clearwater River, Idaho. Pages 169-175 In. Catch-and-release fishing as a management tool. Humboldt State University, Arcata, California.
- Ricker, W.E. 1949. Effects of removal of fins upon the growth and survival of spiny-rayed fishes. Journal of Wildlife Management 13(1):29-40.
- Shetter, D.S. 1951. The mortality and growth of marked and unmarked lake trout fingerlings in the presence of predators. Transactions of the American Fisheries Society 81:17-34.
- Shetter, D.S. and L.N. Allison. 1955. Comparison of mortality between fly-hooked and worm-hooked trout in Michigan streams. Michigan Department of Conservation, Institute of Fisheries Research, Miscellaneous Publication No. 9.
- Stauffer, T.M. and M.J. Hansen. 1969. Mark retention, survival, and growth of jaw-tagged and fin-clipped rainbow trout. Transactions of the American Fisheries Society 98(2):225-229.
- Stuart, T.A. 1957. The migrations and homing behaviour of brown trout (Salmo trutta L.). Scientific Investigation Freshwater Fish. \_\_\_\_\_Scotland 18:27.
- Stuart, T.A. 1958. Marking and regeneration of fins freshwater and salmon. Fishery Research, Scottish Home Department 22:14.
- Wales, J.H. 1947. Growth rate and fin regeneration in trout. Progressive Fish-Culturist 9(2):86-89.
- Weisel, G.F. 1968. Salmonid adipose fin. Copeia. pp. 626-627.



## A P P E N D I C E S

## Appendix A. BIBLIOGRAPHY - Marked Fish Survival

- Armstrong, G. C. 1947. Mortality, rate of growth, and fin regeneration of marked and unmarked lake trout fingerlings at the Provincial Fish Hatchery, Port Arthur, Ontario. Transactions of the American Society. 77: 129-131.
- Barnaby, J. T. 1944. Fluctuations in abundance of red salmon, Oncorhynchus nerka (Walbaum), of the Karluk River, Alaska. Fishery Bulletin, United States Fish and Wildlife Service. 50(39):237-295.
- Bonham, K., and L. R. Donaldson. 1966. Low level chronic irradiation of salmon eggs and alevins. International Atomic Energy Agency.
- Bonham, K. 1968. Growth inhibition by maxillary excision, sized at start of jaw elongation, and regression of *weight* on length of precocious male chinook salmon. Transactions of the American Fisheries Society. 97(2):198-201.
- Brynildson, D. M., and C. L. Brynildson. 1967. The effect of pectoral and ventral fin removal on survival and growth of wild brown trout in a Wisconsin stream. Transactions of the American Fisheries Society. 96(3):353-355.
- Calkins, T. P. 1959. The effect of fin removal on the swimming ability of young silver salmon. University of Washington, Fisheries Research Institute, Circular 109.
- Chadwick, H. K. 1966. Fish marking. A. Calhoun, editor. Inland fisheries management. California Department of Fish and Game.
- Churchill, W. S. 1963. The effect of fin removal on survival, growth, and vulnerability to capture of stocked walleye fingerlings. Transactions of the American Fisheries Society. 92(3):298-300.
- Coble, D. W. 1967. Effects of fin-clipping on mortality and growth of yellow perch with a review of similar investigations. Journal of Wildlife Management. 31(1):173-180.

- Coble, D. U. 1972. Vulnerability of fin clipped bluegill to largemouth bass predation in tanks. Transactions of the American Fisheries Society. 111(3):563-565.
- Cope, O. B. and D. U. Slater. 1957. Role of Coleman Hatchery in maintaining a king salmon run. United States Fish and Wildlife Service Research Report. 47:22.
- Crawford, R. U. 1958. Behavior, growth and mortality in the bluegill, Lepomis macrochirus Rafinesque, following fin clipping. Copela (4) :330 -331.
- Dill, U. A., editor. 1972. Report of the symposium on the major communicable fish disease in Europe and their control Fisheries Assistant Operation Technical Paper No. 17. Rome.
- Eipper, A. W., and J. L. Forney. 1965. Evaluation of partial fin clips for marking largemouth bass, walleyes, and rainbow trout. New York Fish and Game Journal. 12(2):223-240.
- Foerster, R. E. 1936. The return from the sea of sockeye salmon (Oncorhynchus nerka) with special reference to percentage survival, sex proportions and progress of migration. Journal of the Biological Board of Canada. 3(1):26-42.
- Foerster, R. E. 1936. An investigation of the life history and propagation of the sockeye salmon (Oncorhynchus nerka) at Cultus Lake, British Columbia. No. 5. The life history cycle of the 1926 year class with artificial propagation involving the liberation of free-swimming fry. Journal of the Biological Board of Canada. 2(3):211-333.
- Fry, D. H., Jr. 1961. Some problems in the marking of salmonida. Pacific Marine Fisheries Commission Bulletin 5:77-83.
- Harris, J. E. 1936. The role of the fins in the equilibrium of the swimming fish. I. Wind tunnel tests on a model of Mustelus canis (Mitchill). Journal of Experimental Biology. 13:476-493.
- Harris, J. E. 1937. The mechanical significance of the position and movements of the paired fins in the Teleostei Papers from Tortugas Lab. Carnegie Institution of Washington. 31(7):173-189.

- Harris, J. E. 1938. The role of the fins in the equilibrium of the swimming fish. II. The role of the pelvic fins. *Journal of Experimental Biology* 15:32-47.
- Hatch, R. U., and D. A. Webster. 1961. Trout production in four central Adirondack Mountain lakes. Cornell University Agricultural Experimental Station.
- Herman, E. F. 1946. Notes on the effect of cauterization in fin-clipping trout. *Copeia*. Issue 2.
- Herman, R. L. 1968. Fish furunculosis 1952-1966. *Transactions of the American Fisheries Society*. 97(3):221-231.
- Horack, D. C. 1969. The effect of fin removal on stamina of hatchery-reared rainbow trout. *Progressive Fish-Culturist*. 31(4):217-220.
- Houston, A. H., J. A. Madden, R. J. Woods, and If. M. Milen. 1971. Some physiological effects of handling and tricaine methane sulphonate anesthetization upon the brook trout ' (Salvelinus fontinalis). *Journal of Fisheries Research Board of Canada*. 28(5):625-633.
- Johnson, H. E., and J. M. Shelton. 1958. Marking Chinook salmon fry. *Progressive Fish-Culturist*. 20:183-186.
- Lux, F. E. 1960. Notes of first-year growth of several species of Minnesota fish. *Progressive Fish-Culturist* 22(2):81-82.
- Marking, L. L. 1967. Toxicity of MS-222 to selected fishes. United States Fish and wildlife Service, Investigation of Fish Control. 12:1-10.
- Mears, H. C., and R. U. Hatch. 1976. Overwinter survival of fingerling brook trout with single and multiple fin clips. *Transactions of the American Fisheries Society*. 105(6):669-674.
- Mears, H. C. 1976. Effect of fin excision on survival of fingerling brook trout (Salvelinus fontinalis) in a reclaimed pond. Master's Thesis. University of Maine. Orono.
- Nelson, U. C. 1960. A comparison of the effects of removal of the adipose and pelvic fin on fingerling brook and rainbow trout. Colorado Department of Game and Fish.

- Nicola, S. J., and A. J. Cordone. 1973. Effects of fin removal on survival and growth of rainbow trout (Salmo gairdneri) in a natural environment. Transactions of the American Fisheries Society. 102(4):753-759.
- Nielson, R. S., N. Reimers, and H. D. Kennedy. 1957. A six-year study of the survival and vitality of hatchery-reared trout of catchable size in Convict Creek, California. California Fish and Game Bulletin, 43:5-42.
- Parker, R. R., E. C. Black and P. A. Larkin. 1961. Some aspects of fish-marking mortality. International Commission for the Northwest Atlantic Fisheries, North Atlantic Fish Marking Symposium, Woods Hole, Massachusetts. May 1961. Serial No. 828 (B. Rept. 1), Contribution No. 19.
- Parker, R. R., E. C. Black, and P. A. Larkin. 1963. Some aspects of fish-marking mortality. North Atlantic Fish Marking Symposium International Commission. Northwest Atlantic Fisheries Special Publication No. 4:117-122.
- Patrick, B., and R. Haas. 1971. Fins pulling as a technique for marking muskellunge fingerlings. Progressive Fish-Culturist. 33(2):116-118.
- Phillips, A. M., Jr., H. A. Podoliak, D. R. Brockway, and R. R. Vaughn. 1957. The nutrition of trout. Cortland Hatchery Report 26, New York Conservation Department of Fisheries Research Bulletin 21.
- Phinney, D. E., and S. B. Mathews. 1969. Field test, of fluorescent pigment marking and fin-clipping of coho salmon. Journal of Fisheries Research Board of Canada. 26(6):1619-1624.
- Pritchard, A. L. 1939. Homing tendency and age of maturity of pink salmon (Oncorhynchus gorbuscha) in British Columbia. Journal of Fisheries Research Board of Canada. 4(4):233-251.
- Pritchard, A. L. 1941. The recovery of marked Masset Inlet pink salmon during the season of 1940. Fisheries Research Board of Canada, Program Report Pacific Coast Stations. 48:13-17.
- Radcliffe, R. U. 1950. The effect of fin-clipping on the cruising speed of goldfish and coho salmon fry. Journal of Fisheries Research Board of Canada. 8(2):67-73.

- Ricker, W. E. 1945. Abundance, exploitation and mortality of the fishes in two lakes. Investigation of Indiana Lakes and Streams. 2(17):345-448.
- Ricker, W. E. 1948. Effects of removal of fins upon the growth and survival of spiny-rayed fishes. Journal of Wildlife Management. 13:29-40.
- Ricker, W. E. 1949. Effects of removal of fins upon the growth and survival of spiny-rayed fishes. Journal of Wildlife Management. 13(1):29-40.
- Robson, D. .S., and H. A. Realer. 1968. Estimation of population number and mortality rates, Chap. 6 in W.E. Ricker, editor. Methods for assessment of fish production in fresh waters. International Biological Programme Handbook No. 3. Blackwell Scientific Publications, Oxford and Edinburgh.
- Salo, E. O. and U. H. Bayliff. 1958. Artificial and natural production of silver salmon (Oncorhynchus kisutch) at Minter Creek, Washington. Washington Department of Fisheries Research Bulletin 4:1-82.
- Saunders, R. L., and K. R. Allen. 1967. Effects of tagging and of fin-clipping on the survival and growth of Atlantic salmon between smolt and adult stages. Journal of Fisheries Research Board of Canada. 24(12):2595-2611.
- Schoettger, R. A., and A. M. Julin. 1967. Efficacy of MS-222 as an anesthetic: on four salmonids. United States Fish and Wildlife Service. Investigation of Fish, Control 13. Resource Publication 19:1-15.
- Scott, B. 1968. Methods for assessment of fish production in fresh waters. International Biological Programme Handbook No. 3. Blackwell Scientific Publications, Oxford and Edinburgh.
- Senn, H. 1970. Evaluation of 1965 brood coho released from ten Puget Sound and three coastal hatcheries. Washington, Department of Fisheries, Hatcheries Division.
- Shapovalov, L. and A. C. Taft. 1954. The life histories of the steelhead rainbow trout (Salmo gairdneri gairdneri) and silver salmon (Oncorhynchus kisutch) with special reference to Waddell Creek, California, and recommendations regarding their management. California Department of Fish and Game, Fisheries Bulletin 98:375.

- Shetter, D. S. 1951. The effect of fin removal on fingerling lake trout (Cristivomer namaycush). Transactions of the American Fisheries Society. 80:260-277.
- Shetter, D. S. 1952. The mortality and growth of marked and unmarked lake trout fingerlings in the presence of predators. Transactions of the American Fisheries Society. 81:1734.
- Shetter, David S. 1967. Effects of jaw tags and fin excision upon the growth, survival, and exploitation of hatchery rainbow trout fingerlings in Michigan. Transactions of the American Fisheries Society. 96(4):394-399.
- Stauffer, T. M., and M. J. Hanson. 1969. Mark retention, and survival, and growth of jaw-tagged and fin-clipped rainbow trout. Transactions of the American Fisheries Society. 98(2):255-229.
- Stolte, L. U. 1973. Differences in survival and growth of marked and unmarked coho salmon. Progressive Fish-Culturist. 35(4):229-230.
- Stuart, T. A. 1958. Marking and regeneration of fins freshwater and salmon. Fishery Research, Scottish Home Department. 22:14.
- Wales, J. H. 1947. Growth rate and fin regeneration in trout. Progressive Fish-Culturist. 9(2):86-89.
- Wales, J. H., and E. R. German. 1956. Castle Lake investigation-second phase: eastern brook trout. California Fish and Game Bulletin 42(2):93-108.
- Weber, D., and R. G. Wahle. 1969. Effect of fin-clipping on survival of sockeye salmon (Oncorhynchus nerka). Journal of Fisheries Research Board of Canada. 26(5):1263-1271.
- Wolf, L. E. 1954. Development of resistant strains of fish. Transactions of the American Fisheries Society. 83:342-349.

## Appendix B. BIBLIOGRAPHY - Fin Regeneration

- Armstrong, G. C. 1947. Mortality, rate of growth, and fin regeneration of marked and unmarked lake trout fingerlings at the Provincial Fish Hatchery, Port Arthur, Ontario. Transactions of the American Fisheries Society. 77:129-131.
- Bonham, K., and L. R. Donaldson. 1966. Low-level chronic irradiation of saloon eggs and alevins. Disposal of Radioactive Wastes into Seas, Oceans and Surface Waters. International Atomic Energy Agency.
- Bonham, K. 1968. Growth inhibition by maxillary excision, size at start of jaw elongation, and regression of weight on length of precocious male chinook salmon. Transactions of the American Fisheries Society. 97(2):198-201.
- Brynildson, O. M., and C. L. Brynildson. 1967. The effect of pectoral and ventral fin removal on survival and growth of wild brown trout in a Wisconsin stream. Transactions of the American Fisheries Society. 96(3):353-355.
- Calkins, T. P. 1959. The effect of fin removal on the swimming ability of young silver salmon. University of Washington, Fisheries Research Institute, Circular 109.
- Carlander, K. D. 1953. Handbook of freshwater fishery biology. William C. Brown Company, Dubuque, Iowa.
- Chadwick, H. K. 1966. Fish marking. Inland Fisheries Management. California Department of Fish and Game.
- Coble, D. W. 1971. Effects of fin clipping and other factors on survival and growth of smallmouth bass. Transactions of the American Fisheries Society. 100(3):460-473.
- Day, F. 1887. British and Irish Salmonidae. London: Williams and Norgate.
- Eipper, A. U., and J. L. Forney. 1965. Evaluation of the partial fin clips for marking largemouth bass, walleyes, and rainbow trout. New York Fish and Game Journal. 12(2):233-240.*
- Foerster, R. E. 1934. An investigation of the life history and propagation of the sockeye salmon (Oncorhynchus nerka) at Cultus Lake, British Columbia. No. 4. The life history cycle of the 1925 year class with natural propagation.



- Foerster, R. E. 1936. An investigation of the life history and propagation of the sockeye salmon (Oncorhynchus nerka) at Cultus Lake, British Columbia. No. 5. The life history cycle of the 1926 year class with artificial propagation involving the liberation of free-swimming fry. Journal of the Biological Board of Canada. 2(3):26-42.
- Foerster, R. E. 1936. The return from the sea of sockeye salmon (Oncorhynchus nerka) with special reference to percentage survival, sex proportions and progress of migration. Journal of the Biological Board of Canada. 3(1):26-42.
- Fry, D. H., Jr. 1961. Some problems in the marking of salmonids. Pacific Marine Fisheries Commission Bulletin. 5:77-83.
- Hale, J. C. 1954. Regeneration of pelvic fins on fingerling brook trout following fin clipping. Minnesota Department of Conservation, Investigation Report \*149.
- Hallock, R. J., G. H. Warner, and D. H. Fry, Jr. 1952. California's part in a three-state salmon fingerling marking program. California Fish and Game. 38(3):301-332.
- Harris, J. E. 1936. The role of the fins in the equilibrium of the swimming fish. I. Wind tunnel tests on a model of Mustelus canis (Mitchill). Journal of Experimental Biology. 13:476-493.
- Harris, J. E. 1937. The mechanical significance of the position and movements of the paired fins in the Teleostei. Papers from Tortugas Lab. Carnegie Institution of Washington. 31(7):173-189.
- Harris, J. E. 1938. The role of the fins in the equilibrium of the swimming fish. II. The role of the pelvic fins. Journal of Experimental Biology. 15:32-47.
- Hartt, A. C. 1959. Movement of salmon in the North Pacific Ocean and Bering Sea as determined by tagging. University of Washington, College of Fisheries, Circular 106.
- Hartt, A. C. 1962. Movement of salmon in the North Pacific Ocean and Bering Sea as determined by tagging. Bulletin of the International North Pacific Fisheries Commission. 6:1-157.

- Heacox, C. 1942. Fin regeneration in brown trout. Transactions Of the American Fisheries Society. 72:231-232.
- Henry, K. A. 1971. Estimates of maturation and ocean mortality for Columbia River hatchery fall chinook salmon and effect of no ocean fishery on yield. Oregon Fisheries Commission Research Report. 3:13-27.
- Herman, E. F. 1946. Notes on the effect of cauterization in fin-clipped trout. Copeia.
- Horack, D. C. 1969. The effect of fin removal on stamina of hatchery-reared rainbow trout. Progressive Fish-Culturist. 31(4):217-220.
- Hunter, J. G. 1959. Survival and production of pink and chum salmon in a coastal stream. Journal of the Fisheries Research Board of Canada. 16(6):5-886.
- Johnson, E. H., and J. M. Shelton. 1958. Marking chinook salmon fry. Progressive Fish-Culturist. 20(4):183-185.
- Jones, J. U., and J. N. Ball. 1954. The spawning behaviour of brown trout and salmon. British Journal of Animal Behaviour. 2:103-114.
- Lander, R. H., and K. A. Henry. 1973. Survival, maturity, abundance and marine distribution of 1965-66 brood coho salmon. Oncorhynchus kisutch, from Columbia River hatcheries. United States National Marine Fisheries Service, Fisheries Bulletin 71:679-695.
- Lander, R. H. 1973. Problem of bias in models to approximate ocean mortality, maturity and abundance schedules of salmon from known emolts and returns. United States National Marine Fisheries Service, Fisheries Bulletin 71:513-525.
- Larkin, P. A., and J. G. McDonald. 1968. Factors in the population biology of the sockeye salmon of the Skeena River. Journal of Animal Ecology. 37:229-258.
- Lux, F. E. 1960. Notes on first-year growth of several species of Minnesota fish. Progressive Fish-Culturist 22(2):81-82.
- Mears, H. C. 1976. Effect of fin excision on survival of fingerling brook trout (Salvelinus fontinalis) in a reclaimed pond. Master's Thesis. University of Maine.

- Mears, H. C. 1976. Overwinter regeneration of clipped fins in fingerling brook trout. *Progressive Fish-Culturist*. 38(2):73.
- Nelson, Wesley, C. 1960. A comparison of the effects of the removal of the adipose and pelvic fins on fingerling brook and rainbow trout. Colorado Department of Game and Fish.
- Nielson, R. S., N. Reimeers, and H. D. Kennedy. 1957. A six-year study of hatchery-reared trout of catchable size in Convict Creek, California. *California Fish and Game Bulletin* 43(1):5-42.
- Parker, R. R., E. C. Black, and P. A. Larkin. 1963. Some aspects of fish-marking mortality. *North Atlantic Fish Marking Symposium, International Commission North West Atlantic Fisheries Special Publication*. 4:117-122.
- Patrick, B., and R. Haas. 1971. Fin pulling as a technique for marking muskellunge fingerlings. *Progressive Fish-Culturist*. 33(2):116-118.
- Phinney, D. E., and S. B. Mathews. 1969. Field test of fluorescent pigment marking and fin-clipping of coho salmon. *Journal of the Fisheries Research Board of Canada*. 26(6):1619-1624.
- Radcliffe, R. U. 1950. The effect of fin-clipping on the cruising speed of goldfish and coho salmon fry. *Journal of the Fisheries Research Board of Canada*. 8(2):67-73.
- Rich, U. H., and H. B. Holmes. 1928. Experiments in marking young chinook salmon on the Columbia River, 1916 to 1927. *Bulletin of the United States Bureau of Fisheries*. 44:215-264.
- Ricker, W. E. 1946. Abundance, exploitation and mortality of the fishes in two lakes. *Investigations of Indiana lakes and streams*. 2(17):345-448.
- Ricker, W. E. 1949. Effects of removal of fins upon the growth and survival of spiny-rayed fishes. *Journal of Wildlife Management*. 13(1):29-40.
- Ricker, W. E. 1958. Handbook of computation for biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada* 119.

- Robson, D. S., and H. A. Regier. 1968. Estimation of population number and mortality rates. Methods for assessment of fish production in fresh waters. International Biological Programme Handbook No. 3. Blackwell Scientific Publications, Oxford and Edinburgh.
- Saunders, R. L., and K. R. Allen. 1967. Effects of tagging and of fin-clipping on the survival and growth of Atlantic salmon between smolt and adult stages. Journal of the Fisheries Research Board of Canada. 24(12):2595-2611.
- Scott, B. 1968. Marking and tagging. Methods for assessment of fish production in fresh waters. International Biological Programme Handbook No. 3. Blackwell Scientific Publications, Oxford and Edinburgh.
- Shatter, D. S. 1950. The effect of fin removal on Fingerling lake trout (Cristivomer namaycush). Transactions of the American Fisheries Society. 80:260-277.
- Shatter, D. S. 1951. The mortality and growth of marked and unmarked lake trout fingerlings in the presence of predators. Transactions of the American Fisheries Society. 81:17-34.
- Shetter, D. S. 1967. Effects of jaw tags and fin excision upon the growth, survival and exploitation of hatchery rainbow trout fingerlings in Michigan. Transactions of the American Fisheries Society. 96(4):394-399.
- Slater, D. U. 1949. Re-formation of excised fins of king salmon fingerlings and its effects on recognition of marked adults. Transactions of the American Fisheries Society. 77:132-140.
- Stauffer, T. M., and M. J. Hansen. 1969. Mark retention, survival, and growth of jaw-tagged and fin-clipped rainbow trout. Transactions of the American Fisheries Society. 98(2):225-229.
- Stolte, L. U. 1973. Differences in survival and growth of marked and unmarked coho salmon. Progressive Fish-Culturist. 35(4):229-230.
- Stuart, T. A. 1953. Spawning migration, reproduction and young stages of loch trout (Salmo trutta L.). Scientific Investigation of Freshwater Fish. 5:39.

- Stuart, T. A. 1957 The migrations and homing behaviour of brown trout (Salmo trutta L.) Scientific Investigation Freshwater Fish. 18:27.
- Stuart, T. A. 1958. Marking and regeneration of fins. Freshwater and Salmon Fisheries Research, Scottish Home Department. 22:14.
- Taguchi, K. 1961. A trial to estimate the instantaneous, rate of natural mortality of adult salmon (Oncorhynchus sp.) and the consideration of rationality of offshore fishing. I. For chum salmon (Oncorhynchus keta). Bulletin of the Japanese Society of Science and Fish. 27:963-971.
- Wales, J. H. 1947. Growth rate and fin regeneration in trout. Progressive Fish-Culturist. 9(2):86-89.
- Wales, J. H., and E. R. German. 1956. Castle Lake investigation-second phase: eastern brook trout. California Fish and Game Bulletin 42(2):93-108.
- Worlund, D. D., R. J. Walle, and P. D. Zimmer. 1969. Contribution of Columbia River hatcheries to harvest of fall Chinook salmon (Oncorhynchus tshawytscha). United States Fish and Wildlife Service, Fishery Bulletin 67:361-391.

## Appendix C. BIBLIOGRAPHY - Hookinej Mortality

- Adams, J. R. 1960. A catchable trout study in the Truckee River. Master's Thesis, Humboldt State College, Arcata California.
- Alexander, R. 1974. Functional design in fishes. Hutchinson and Company. London, England.
- Anon. 1969. A summary or information of the "shaker" catch by Washington's ocean salmon fisheries. Washington Department of Fish.
- Ball, K. W. 1971. Initial effects of catch-and-release regulations on cutthroat trout in an Idaho stream. Master's Thesis, University of Idaho, Moscow, Idaho.
- Barnhart, R, A., and T. D. Roelofs, editors. 1977. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Benson, N. G., and R. V. Bulkley. 1963. Equilibrium yield and management of cutthroat trout in Yellowstone Lake. United States Fish and Wildlife Service, Research Report 62:44.
- Beukema, J. J. 1969. Angling experiments with care (Cypinus carpio L.). I. Differences between wild, domesticated, and hybrid strains. Netherlands Journal of Zoology. 19(4):596-609.
- Beukema, J. J. 1970. Acquired hook-avoidance in the pike (Esox lucius L.) fished with artificial and natural baits. Journal of Fishery Biology. 2(2):155-160.
- Bjornn, T. C . 1969. Hatchery cutthroat trout, hooking mortality study. Unpublished data. Idaho Cooperative Fishery Research Unit, University of Idaho, Moscow, Idaho.
- Bjornn, T. C., T. H. Johnson, and R. F. Thurow, 1977. Angling versus natural mortality in northern id d l w cutthroat trout populations. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Black, E. C. 1958. Hyperactivity as a lethal factor in fish. Journal of the Fisheries Research Board of Canada. 15:573-586.

- Bouck, G. R., and R. C. Ball. 1966. Influence of capture methods on blood characteristics and mortality in the rainbow trout (Salmo gairdneri). Transactions of the American Fisheries Society. 95(2):170-176.
- Brett, J. R. 1958, implications and assessments of environmental stress. The investigation of fish-power problems. H. R. MacMillan Lectures in Fisheries, University of British Columbia, Vancouver.
- Buchanan, C. C., F. J. Mather, and J. C. Mason, Jr. 1977. An overview: Cooperative game fish tagging program (Atlantic Ocean). Catch-and-release fishing as a Management tool. Humboldt State University, California.
- Burns, D. C. 1977, Catch-and-release fishing for steelhead in the State of Washington. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Butler, R.L., and D. P. Borgenson. California "catchable" trout fisheries. Fisheries Bulletin 127, California Department of Fish and Game.
- Calkins, T. P. 1959. The effect of fin removal on the swimming ability of young silver salmon. Fisheries Research Institute, University of Washington. Circular 109:6.
- Carpenter, R. M., and J. M. Deinstadt. 1977. Attitudes of trout stream anglers and their relation to trout stream management. Catch--and-release fishing as a management tool. Humboldt State University, California.
- Cartwright, J. W. 1961. Hooking mortality study - caribou region - 1961. British Columbia Fish and Wildlife Branch, Management Report 50.
- Casillas, E., and L. S. Smith. 1977. Effect of stress on blood coagulation and hematology in rainbow trout (Salmo gairdneri). Journal of Fisheries Biology. 10(5) :481-491.
- Churchill, W. S. 1963. The effect of fin removal on survival, growth, and vulnerability to capture of stocked walleye fingerlings. Transactions of the American Fisheries Society. 92:298-300.
- Cleaver, F. C. 1969. Effects of ocean fishing on 1961-brood fall Chinook salmon from Columbia River hatcheries. Oregon Fisheries Commission Research Report. 1:76.

- Cochran, W. G. 1977. Sampling techniques. Wiley and Sons, New York.
- Cooksey, R. W., and B. L. Driver. 1977. Preferred psychological outcomes of recreational fishing. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Cooper, E. L. 1952. Returns from plantings of legalized brook, brown and rainbow trout in the Pigeon River, Otsego County, Michigan, Transactions of the American Fisheries Society. 82:265-280.
- Cordes, R. A. 1977. The fly fisherman's view of catch-and-release fishing. Catch--anal-release fishing as a management tool. Humboldt State University, California.
- Cresswell, R. C. 1981. Post-stocking movements and recapture of hatchery-reared trout released into flowing waters - a review. Journal of Fisheries Biology. 18:429-442.
- Deinstadt, J. M. 1977. Catch-and-release angling in California's wild trout streams. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Dotson, T. 1982. Mortalities in trout caused by gear type and angler-induced stress. North American Journal of Fisheries Management 2:60-65.
- Falk, M. R., D. V. Gillman, and L. N. Dahlke. 1974. Comparison of mortality between barbed and barbless hooked lake trout. Environment Canada, Fisheries Marine Service, Resource Management Branch, Central Region, Technical Report Series No. CEN/T-74-1.
- Falk, M. R., and D. V. Gillman. 1975. Mortality data for angled arctic grayling and northern pike from the great Slave Lake area, Northwest Territories. Environment Canada, Fisheries Marine Service, Resource Management Branch, Central Region, Data Report Series No. CEN/D-75-1.
- Gowing, H. 1975. Population dynamics of wild brown trout in Gamble Creek, subject first to angling, then no angling. Fisheries Research Report No. 1824. Michigan Department of Natural Resources.
- Graff, D. R. and B. A. Hollander. 1977. Catch-and-release fishing -- the Pennsylvania experience. Catch-and-release fishing as a management tool. Humboldt State University, California.



- Grasswell, R. E. 1976. Hooking mortality in trout. United States Department of Interior, National Park Service, Yellowstone National Park, Informational Paper No. 29.
- Griggs, J. D. 1977. Roemer Lake -- Atlantic salmon catch-and-release program. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Guataveson, A. U. 1977. Physiological response to hooking stress in largemouth bass, Micropterus salmoides. Master's Thesis, Utah State University, Logan, Utah.
- Harris, J. E. 1936. The role of the fins in the equilibrium of the swimming fish. I. Wind tunnel tests on a model of Mustelus canis (Mitchill). Journal of Experimental Biology. 13:476-493.
- Harris, J. E. 1937. The mechanical significance of the position and movements of the paired fins in the Teleostei. Papers from Tortugas Lab. Carnegie Institution of Washington. 31(7):173-189.
- Harris, J. E. 1938. The role of the fins in the equilibrium of the swimming fish. It. The role of the pelvic fine. Journal of Experimental Biology. 15:32-47.
- Hattingh, J., and A. J. J. VanPletzen. 1974. The influence of capture and transportation on some blood parameters of freshwater fish. Comparative Biochemistry Physiology. 49A:607-609.
- Haw, F. 1963. Comparative hooking mortality between treble and single hooks on Chinook salmon. Pacific Marine Fisheries Commission, Annual Report 15:22-24.
- Helmer, J. T., W. M. Frazier, and J. S. Griffith. 1985. Post-stocking performance of catchable-size hatchery rainbow trout with and without pectoral fins. North American Journal of Fisheries Management. 5:21-25.
- Helrich, L. A., and U. T. Kendall. 1982. Movements of hatchery-reared rainbow, brook and brown trout stocked in a Virginia mountain stream. The Progressive Fish-Culturist. 44:3-7.
- Hollander, B. A., and D. R. Graff. 1977. Catch-and-release fishing - the Pennsylvania experience. Catch-and-release fishing as a management tool. Humboldt State University, California.

- Horack, D. L., and W. D. Klein. 1967. Influence of capture methods on fishing success, stamina, and mortality of rainbow trout in Colorado. Transactions of the American Fisheries Society. 96(2):220-222.
- Hunsaker, D., II, L. F. Marnell, and F. P. Sharpe. 1970. Hooking mortality of Yellowstone cutthroat trout. Progressive Fish-Culturist. 32(4):231-235.
- Hunt, R. L., O. M. Brynildson, and J. T. McFadden. 1962. Effects of angling regulations on a wild brook trout fishery. Wisconsin Department of Natural Resources, Technical Bulletin No. 26.
- Hunt, R. L. 1975. Angling regulations in relation to wild trout management. Wild Trout Management, Trout Unlimited Incorporated, Denver, Colorado.
- Hunt, R. L. 1977. An unsuccessful use of catch-and-release regulations for a wild brook trout fishery. Catch-and-release fishing as a management tool, Humboldt State University, California.
- Huntsman, A. G. 1938. Overexertion as a cause of death of captured fish. Science. 87(2269):577-578.
- Hurn, D. R. 1961. Gold River hooking mortality on summer steelhead. Unpublished data. British Columbia Fish and Wildlife Branch.
- Johnson, T. H., and T. C. Bjornn. 1975. Evaluation of angling regulations in management of cutthroat trout. Job Performance Report: F-59-R-6. University of Idaho, Cooperative Fisheries Research Unit, Moscow, Idaho.
- Johnson, T. H. 1977. Catch-and-release and trophy-fish regulations in the management of cutthroat trout populations and fisheries in Northern Idaho streams. Master's Thesis. University of Idaho, Moscow, Idaho.
- Klein, W. D. 1969. The use of special regulations in trout stream management in Colorado. Colorado Game, Fish and Parks Division.
- Klein, W. D. 1966. Mortality of trout caught on artificial lures and released by fishermen. Transactions of the American Fisheries Society. 95(3):326-328.

- Klein, U. D. 1974. Mortality among trout released by anglers while ice-fishing. Unpublished Master's Thesis, Colorado State University. Colorado Division of Wildlife, Fort Collins.
- Lasater, J. E., and F. Haw. 1961. Comparative hooking mortality between treble and single hooks on silver salmon. Pacific Marine Fisheries Commission. 5:73-76.
- Latta, U. C. 1973. The effects of a flies-only fishing regulation upon trout in the Pigeon River, Otsego County, Michigan. Michigan Department of Natural Resources, Fisheries Division, Fisheries Research Report No. 1807.
- Lennon, R. E. and P. S. Parker. 1960. The fishing-for-fun program on trout streams in Great Smoky Mountains National Park. Proceedings of the Society of the American Foresters. 1960:106-112.
- MacPhee, C. 1966. Influence of differential angling mortality and stream gradient on fish abundance in a trout-sculpin biotope. Transactions of the American Fisheries Society. 95:381-387.
- Marnell, L. F., and D. Hunsaker, II. 1970. Hooking mortality of lure-caught cutthroat trout (Salmo clarki) in relation to water temperature, fatigue, and reproductive maturity of released fish. Transactions of the American Fisheries Society. 99(4):684-688.
- May, B. E. 1972. Evaluation of large scale release programs with special reference to bass fishing tournaments. Proceedings of the 26th Annual Conference Southeastern Association Game and Fish Commission. 26:325-329.
- Mazeaud, M. M., F. Mazeaud, and E. M. Donaldson. 1977. Primary and secondary effects of stress in fish: Some new data with a general review. Transactions of the American Fisheries Society. 106:201-212.
- McLaren, J., and R. L. Butler. 1970. The Spruce Creek catch-and-release study. Trout Unlimited, 11:18-19, 29.
- McLaren, J., and R. L. Butler. 1970. Vulnerability of wild brown trout to fly fishing under catch-and-release regulations. Master's Thesis, Pennsylvania State University.

- Milne, D. J., and E. A. R. Ball. 1956. The mortality of small salmon when caught trolling and tagged or released untagged. Fisheries Research Board of Canada Program Report 106:10-13.
- Mongillo, P. E. 1976. A bioenergetic study of brown trout in a natural stream. Master's Thesis. Utah State University.
- Moody, H. L. 1974. Tournament catch of largemouth bass from St. Johns River, Florida. Proceedings of the 28th Annual Conference of the Southeastern Association Game and Fish Commission. 28:73-82.
- Newell, A. E. 1957. Two-year study of movements of stocked brook trout and rainbow trout in a mountain trout stream. The Progressive Fish-Culturist. 19:76-80.
- Ortmann, D. J. 1976. Steelheading-just for fun? Idaho Wildlife Review.29(1):8-10.
- Parker, R. R., and E. C. Black. 1959, Muscular fatigue and mortality in troll-caught chinook salmon (Oncorhynchus tshawytscha) Journal of the Fisheries Research Board of Canada. 16:95-106.
- Parker, R. R., E. C. Black, and P. A. Larkin. 1959. Fatigue and mortality in troll-caught Pacific salmon. Journal of the Fisheries Research Board of Canada. 16:429--448.
- Peltzman, R. J. 1978. Hooking mortality of juvenile largemouth bass, Micropterus salmoides. California Fish and Game Bulletin 64(3):185-188.
- Pettit, S. U. 1977. Steelhead catch-and-release research and regulations on the Clearwater River, Idaho. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Radcliffe, R. W. 1950. The effect of fin-clipping on the cruising speed of goldfish and coho salmon fry. Journal of the Fisheries Research Board of Canada. 8(2):67-73.
- Reed, P. H. 1970. A review of trolling gear studies on chinook and coho salmon. Fisheries Commission of Oregon.

- Rutledge, W. F., and D. L. Pritchard. 1977 Hunting mortality of largemouth bass captured by artificial lures and natural bait. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Seaman E. A. 1969. Quantity vs. quality in fisheries management. Annual Conference Western Association Game and Fish Commission Proceedings. 49:60-67.
- Seidensticker, E. P. 1977. Mortality of largemouth bass for two tournaments utilizing a "Don't Kill Your Catch" program. Catch-and-release fishing as a management tool. Humboldt State University, California.
- Shetter, D. S. 1951. The mortality and growth of marked and unmarked lake trout fingerlings in the presence of predators. Transactions of the American Fisheries Society. 81:17-34.
- Shetter, D. S., and L. N. Allison. 1958. Immortality of trout caused by hooking and artificial lures in Michigan waters, 1956-57. Michigan Department of Conservation, Institute of Fisheries Research, Miscellaneous Publication No. 12.
- Shetter, D. S., and G. R. Alexander. 1962. Effects of flies-only restriction on angling and on fall trout populations in Hunt Creek, Montmorency County, Michigan. Transactions of the American Fisheries Society. 91:295-302.
- Shetter, D. S., and G. R. Alexander. 1966. Angling and trout populations on the North Branch of the Au Sable River, Crawford and Otsego Counties, Michigan, under special and normal regulations, 1958-1963. Transactions of the American Fisheries Society. 95:85-91.
- Shetter, D. S. 1967. Effects of jaw tags and fin excision upon the growth, survival and exploitation of hatchery rainbow trout fingerlings in Michigan. Transactions of the American Fisheries Society 96(4):394-399.
- Squire, J. L. Jr. 1977. The Pacific area cooperative marine game fish tagging program for billfish - and a preliminary determination of striped marlin (Tetrapturus audax) migration patterns in the eastern Pacific. Catch-and-release fishing as a management tool. Humboldt State University, California.

- Varley, J. D. 1976. Yellowstone National Park fishing regulation - biological benefits and liabilities. Annual Conference of the Western Association Game and Fish Commission, Proceedings 56:148-153.
- Wallis, O. L. 1971. Management of aquatic resources and sport fishing in national parks by special regulations. Annual Conference of the Western Association Game and Fish Commission, Proceedings 51:150-163.
- Wardle, C. S. 1972. The changes in blood glucose in Pleuronectes platessa following capture from the wild: a stress reaction. Journal of Marine Biology Association of the United Kingdom. 52(3):635-651.
- Warner, K. 1976. Hooking mortality of landlocked atlantic salmon (Salmo salar) in a hatchery environment. Transactions of the American Fisheries Society. 105(3):365-369.
- Warner, K. 1978. Hooking mortality of lake-dwelling landlocked atlantic salmon (Salmo salar). Transactions of the American Fisheries Society. 107(4):518-522.
- Warner, K., and P. R. Johnson. 1978. Mortality of landlocked atlantic salmon (Salmo salar) hooked on flies and worms in a river nursery area. Transactions of the American Fisheries Society. 107(6):772-775.
- Webster, D. A., and G. S. Little. 1947. Angling experiments by the Fly Tyers' Club at Cornell University. Transactions of the American Fisheries Society. 74(1944):63-70.
- Wedemeyer, G. 1970. The role of stress in the disease resistance of fishes. Symposium on disease of fishes and shellfishes. American Fisheries Society Special Publication No. 5.
- Wedemeyer, G. 1972. Some physiological consequences of handling stress in the juvenile coho salmon (Oncorhynchus kisutch) and steelhead trout (Salmo kisutch). Journal of the Fisheries Research Board of Canada. 29:1730-1738.
- Weithman, A. S., and R. O. Anderson. 1976. Angling vulnerability of esocidae. Annual Conference Southeastern Association Game and Fish Commission, Proceedings 30:99-102.

- Weithman, A. S., and R. O. Anderson. 1977. Catch-and-release fishing for largemouth bass on Hunter's Lake - 1963-1977. Catch-and-release fishing as a management tool Humboldt State University, California.
- Wright, S. 1970. A review of the subject of hooking mortalities in pacific salmon (Oncorhynchus spp.) Washington Department of Fish.
- Wydoski, R. S. 1970. Management of Washington lakes for quality fishing - the Lenice Lake study. Unpublished data. Washington Cooperative Fisheries Research Unit, University of Washington, Seattle.
- Wydoski, R. S., G. A. Wedemeyer, and N. O. Nelson. 1976. Physiological response to hooking stress in hatchery and wild rainbow trout (Salmo gairdneri). Transactions of the American Fisheries Society. 105(5):601-606.
- Wydoski, R. S., and G. A. Wedemeyer. 1976. Problems in the physiological monitoring of wild fish populations. Annual Conference Western Association Game and Fish Commission, Proceedings 56:200-214.

Submitted by:

Rodney C. Duke  
Senior Fishery Research Biologist


Approved by:

IDAHO DEPARTMENT OF FISH AND GAME



---

Jerry M. Conley, Director



---

Steven M. Huffaker, Chief  
Bureau of Fisheries



---

Bill Hutchinson  
Anadromous Hatcheries Supervisor